

ITR/SY: Center for Modeling of Quantum Dynamics, Relaxation and Decoherence in Solid-State Physics for Information-Technology Applications

PI: Vladimir Privman, Clarkson University, DMR Award 0121146


The main **objective** of our program has been to explore *coherent quantum mechanical processes* in novel solid-state semiconductor information processing devices, with components of atomic dimensions. These include quantum computers, spintronic devices, and nanometer-scale logic gates.

Our **achievements** to date include:


New measures of initial decoherence, and evaluation of decoherence for spins in semiconductors.

Evaluation of solid-state quantum computing designs, and studies of transport associated with quantum measurement.

Investigation of spin-polarized devices and role of nuclear spins in spintronics and quantum computing.



Donor electron spin in silicon decoheres owing to interactions with phonons and nuclear spins



Spin-flip scattering in semiconductor nano-device

ITR/SY: Center for Modeling of Quantum Dynamics, Relaxation and Decoherence in Solid-State Physics for Information-Technology Applications

PI: Vladimir Privman, Clarkson University, DMR Award 0121146

Our program has involved four co-PIs: **M.-C. Cheng, M. L. Glasser, D. Mozyrsky, Ch. Tamon**, and extensive collaborations with leading experimental groups and with Los Alamos National Laboratory.

The **educational impact** has included training of 3 undergraduate students, 4 graduate students, 4 postdoctoral researchers, and development of a new course to introduce quantum device concepts to Physics and Electrical & Computer Engineering students.

Our **outreach program** has included sponsoring presentation events, and an international workshop *Quantum Device Technology*, held at Clarkson University in May of 2002.

Our results were published in several papers in refereed journals, and we set up a web site for outreach.


For a spin system in a bosonic bath, we find

$$\|\delta\| = \left[(\rho_{\uparrow\uparrow}(0) - \rho_{\downarrow\downarrow}(0)) \frac{(1 - e^{-B^2(t)})}{2} \right]^2 + \left[(1 - e^{-B^2(t)}) |\rho_{\uparrow\downarrow}(0)| \sin\left(\frac{\Omega}{2}t - \varphi_0\right) \right]^2 =$$


$$= (1 - e^{-B^2(t)}) \left(\frac{(\rho_{\uparrow\uparrow}(0) - \rho_{\downarrow\downarrow}(0))^2}{4} + |\rho_{\uparrow\downarrow}(0)|^2 \sin^2\left(\frac{\Omega}{2}t - \varphi_0\right) \right)^{1/2}$$

where the norm for the density operator deviation is defined as


$$\|\delta\| = \sup_{\varphi \neq 0} \frac{\|(\rho - \rho^{(0)})|\varphi\rangle\|}{\| |\varphi\rangle \|}$$



Transport through NSPI QW



Conductance quantization



Local nuclear spin polarization